

Realistic Decoherence Free Subspaces

K. M. Fonseca Romero,¹ S. G. Mokalzel,² M. O. Terra Cunha,³ and M. C. Nemes^{4,5}

¹*Departamento de Física, Universidad Nacional, Bogotá, Colombia*

²*Departamento de Física, Pontifícia Universidade Católica de São Paulo,
R. Marquês de Paranaguá, 111, São Paulo, S. P., Brazil*

³*Departamento de Matemática, Universidade Federal de Minas Gerais, C.P. 702, Belo Horizonte, 30123-970, Brazil*

⁴*Departamento de Física-Matemática, Instituto de Física,
Universidade de São Paulo, C.P. 66318, 05315-970 São Paulo, S.P., Brazil*

⁵*Departamento de Física, Universidade Federal de Minas Gerais, C.P. 702, Belo Horizonte, 30123-970, Brazil*

(Dated: February 1, 2008)

Decoherence free subspaces (DFS) is a theoretical tool towards experimental implementation of quantum information storage and processing. However, they represent an experimental challenge, since conditions for their existence are very stringent. This work explores the situation in which a system of N oscillators coupled to a bath of harmonic oscillators is close to satisfy the conditions for the existence of DFS. We show, in the Born-Markov limit and for small deviations from separability and degeneracy conditions, that there are *weak decoherence subspaces* which resemble the original notion of DFS.

PACS numbers: 03.65.Yz, 03.67.Lx

I. INTRODUCTION

The very same mechanism responsible for the potential improvements on computation speed using quantum mechanics, is the one which greatly hinders immediate technical implementation. *Entanglement* between different subsystems is essential for the production of the states used in information processing[1]; at the same time, as these qubits can not be completely isolated from its environment, entanglement with the environmental degrees of freedom is a general feature. The deleterious effect of this coupling is usually called *decoherence*[2]. Therefore much effort has been devoted to finding ways around decoherence in quantum computation, such as error correcting codes[3], dynamical decoupling[4] and computation in decoherence free subspaces[5, 6, 7]. Experimental observations of decoherence free evolution have been reported[8, 9]. Many physical implementations have been proposed including cavity QED[10], ions traps[11], nuclear magnetic resonance[12] and semiconductor quantum dots[13]. From the theoretical point of view, recent work has been mainly on proving the existence of DF subspaces, in general related to symmetries of the system which are preserved by the interaction with the environment, on searching for mechanisms of dynamical creation of DFS[14], and on the analysis of their robustness[15]. More realistic models[7] are scarce, and fail to provide insight on the effects of slightly relaxing the conditions necessary for the existence of DFS.

In the present work we consider the case of N independent oscillators linearly coupled to a single environment, and show that the existence of strict decoherence free subspaces can be obtained under the following two conditions: degeneracy of the oscillators, and separability of the coupling with the environment. Both can be viewed as consequences of symmetries: the first involving only the system, and the second the interaction. The exact form of the spectral density and the temperature of the environment are immaterial in what concerns the existence of DFS, since they really decouple from the environment. Master equations for the evolution of the reduced density matrix of the system are derived with and without the Born-Markov approximation, and only the coefficients vary from one case to the other. For two independent oscillators we solve the dynamics of the reduced density and explicit the decoherence free subspace. Also in the case of two harmonic oscillators we study the effect of relaxing the degeneracy and separability conditions. We verify that there is no more DFS, but there remains a long leaved mode, which we call *weak decoherence mode*, and its counterbalance, a *strong decoherence mode* also appear. The time scales for the duration of these components are derived in terms of the appropriate parameters. We analyze these findings in the context of the robustness proof for DFS presented in Ref. [15].

This contribution is organized as follows: in section II an introduction to the concept of decoherence free subspaces is given, in section III the model with oscillators is described, and the decoherence free modes are exhibited. A short discussion of the classical counterpart of decoherence free modes is made. Section IV is devoted to the derivation of master equations. In section V we discuss in details the case of two harmonic oscillators. As a simplifying tool, we introduce the notion of *superoperator*. The section VI discuss the case of small departures from the degeneracy and separability conditions, and shows that despite the fact that the concept of DFS is no more applicable, there remains a *weak decoherence mode* which can be useful for quantum information storing. We give some concluding remarks. Some intermediate calculations have been relegated to the appendix.

II. A QUICK WAY TO DECOHERENCE FREE SUBSPACES

The notion of decoherence free subspaces (DFS) can be easily captured by considering a special coupling between a system and its environment. Consider a system with its autonomous Hamiltonian $\hat{\mathbf{H}}_S$, an environment described by $\hat{\mathbf{H}}_E$, and the interaction between them given by $\hat{\mathbf{H}}_I$. The complete Hamiltonian is then

$$\hat{\mathbf{H}} = \hat{\mathbf{H}}_S + \hat{\mathbf{H}}_E + \hat{\mathbf{H}}_I. \quad (1)$$

Suppose the interaction term can be written in the form $\hat{\mathbf{H}}_I = \hat{\mathbf{A}}_S \hat{\mathbf{B}}_E$, with $\hat{\mathbf{A}}_S$ (resp. $\hat{\mathbf{B}}_E$) acting only on the system (resp. environment) degrees of freedom. We will call this a *separability condition*. Any common eigenvector $|a\rangle$ of $\hat{\mathbf{A}}_S$ and $\hat{\mathbf{H}}_S$ (with eigenvalues a and h_a) does not get entangled with the environment, since in this case

$$\begin{aligned} \hat{\mathbf{H}} |a\rangle \otimes |\epsilon\rangle &= (\hat{\mathbf{H}}_S |a\rangle) \otimes |\epsilon\rangle + |a\rangle \otimes (\hat{\mathbf{H}}_E |\epsilon\rangle) + (\hat{\mathbf{A}}_S |a\rangle) \otimes (\hat{\mathbf{B}}_E |\epsilon\rangle) \\ &= |a\rangle \otimes (h_a + \hat{\mathbf{H}}_E + a \hat{\mathbf{B}}_E) |\epsilon\rangle. \end{aligned} \quad (2)$$

By linearity, any common eigenspace of $\hat{\mathbf{A}}_S$ and $\hat{\mathbf{H}}_S$ is a DFS of the system.

Degeneracy is generally originated by symmetry. Therefore, if one finds a symmetric system where interaction with the environment preserves this symmetry, then any eigenspace of the system is a DFS.

In the language of Ref. [6], $\hat{\mathbf{A}}_S$ is the only *error generator*, and (common) eigenspaces of (all) error generators are DFS. An important distinction that we make is to consider as DFS only the common eigenspaces of $\hat{\mathbf{A}}_S$ and $\hat{\mathbf{H}}_S$, *i.e.*: the system Hamiltonian should not take the state out of a DFS.

III. A MODEL WITH OSCILLATORS

We now present a different situation in which DFS can be achieved. The system will consist of N identical harmonic oscillators (frequency ω , annihilation operators $\hat{\mathbf{a}}_i$, we use $\hbar = 1$). The environment will be modelled as a huge set of harmonic oscillators (frequencies ω_k , annihilation operators $\hat{\mathbf{b}}_k$). Linear coupling will be considered and the rotating wave approximation applied. This model is both: simple enough to be studied in details and general enough to keep the characteristic behaviour of the problem. It is also adequated to make the link with experimental implementations: one can consider vibronic states of N ions trapped together, a system in which (approximate) DFS has already been demonstrated[9, 16], or modes of distinct cavities[17], or even, for $N = 2$, two degenerated modes of one cavity. The Hamiltonian to be considered is

$$\hat{\mathbf{H}} = \omega \sum_{i=1}^N \hat{\mathbf{a}}_i^\dagger \hat{\mathbf{a}}_i + \sum_k \omega_k \hat{\mathbf{b}}_k^\dagger \hat{\mathbf{b}}_k + \sum_{i,k} (g_{ik}^* \hat{\mathbf{a}}_i^\dagger \hat{\mathbf{b}}_k + g_{ik} \hat{\mathbf{a}}_i \hat{\mathbf{b}}_k^\dagger). \quad (3)$$

As in the previous model, we need an additional assumption on the form of the interaction. Assume the coupling constants g_{ik} can be factorized as $G_i D_k$. This can be interpreted as supposing that all oscillators feel the environment in the same way, possibly with only a difference in strength, which depends only on the oscillator itself, not on the environment (the most usual models consider $G_i = G$, which is a special form of the here proposed model). With this factorizability hypothesis, one can rewrite the Hamiltonian (3) as

$$\hat{\mathbf{H}} = \omega \sum_{i=1}^N \hat{\mathbf{a}}_i^\dagger \hat{\mathbf{a}}_i + \sum_k \omega_k \hat{\mathbf{b}}_k^\dagger \hat{\mathbf{b}}_k + \sum_k \left(D_k^* \left(\sum_i G_i^* \hat{\mathbf{a}}_i^\dagger \right) \hat{\mathbf{b}}_k + D_k \left(\sum_i G_i \hat{\mathbf{a}}_i \right) \hat{\mathbf{b}}_k^\dagger \right). \quad (4)$$

By defining the collective operators

$$\hat{\mathbf{A}}_1^\dagger = \frac{\sum_i G_i^* \hat{\mathbf{a}}_i^\dagger}{\sum_i |G_i|^2}, \quad \hat{\mathbf{A}}_1 = \frac{\sum_i G_i \hat{\mathbf{a}}_i}{\sum_i |G_i|^2}, \quad (5)$$

it takes the form

$$\hat{\mathbf{H}} = \omega \sum_{i=1}^N \hat{\mathbf{a}}_i^\dagger \hat{\mathbf{a}}_i + \sum_k \omega_k \hat{\mathbf{b}}_k^\dagger \hat{\mathbf{b}}_k + \hat{\mathbf{A}}_1^\dagger \sum_k c_k \hat{\mathbf{b}}_k + \hat{\mathbf{A}}_1 \sum_k c_k^* \hat{\mathbf{b}}_k^\dagger, \quad (6)$$

where it is clear that only the collective mode $\hat{\mathbf{A}}_1$ is coupled to the environment (in the above formula $c_k = \sum_i |G_i|^2 D_k^*$). One can consider the mode $\hat{\mathbf{A}}_1$ as the first of a new set of normal modes $\{\hat{\mathbf{A}}_i\}$, and the remaining modes thus constitute an infinite dimensional DFS. With this new set of variables, the Hamiltonian is finally written as

$$\hat{\mathbf{H}} = \omega \sum_{i=1}^N \hat{\mathbf{A}}_i^\dagger \hat{\mathbf{A}}_i + \sum_k \omega_k \hat{\mathbf{b}}_k^\dagger \hat{\mathbf{b}}_k + \hat{\mathbf{A}}_1^\dagger \sum_k c_k \hat{\mathbf{b}}_k + \hat{\mathbf{A}}_1 \sum_k c_k^* \hat{\mathbf{b}}_k^\dagger. \quad (7)$$

While the situation in the previous section is completely quantum mechanical, the present model does have a classical analog, because the manipulation above can also be done with classical oscillators. In fact, there is a very old classical situation to which this analysis can be applied: synchronization of pendular clocks. It is known that two clocks in the same wall tend to synchronize in anti-phase. Each clock can be considered as an oscillator, and their coupling to the environment can be considered in terms of the two normal modes (phase and anti-phase modes). The in phase mode couples (much more) strongly to the environment, and this causes the synchronization.

Another consistent analogy of the above model is with the superradiance in the Dicke model[18]. In this case, N two-level atoms are coupled to one field mode. In the regime in which the atoms are collectively coupled to the field (*i.e.*: the field does not distinguish which atom has emitted), the radiative process can be stronger compared to individual emissions (due to interference). The counterpart of this process is the subradiance, *i.e.*: other collective states with weaker emission than the individual contributions (destructive interference). DFS can thus be compared to subradiant states of the system.

IV. MASTER EQUATION

As one usually does not have control on the environment degrees of freedom, the natural approach to this problem is to study the reduced dynamics of the N oscillators. A long but straightforward procedure[19] can be applied to derive the master equation

$$\begin{aligned} \frac{d\hat{\rho}}{dt} = & \frac{1}{i\hbar} [\hat{\mathbf{H}}_0, \hat{\rho}] + (\lambda + \epsilon) \left(2\hat{\mathbf{A}}_1 \hat{\rho} \hat{\mathbf{A}}_1^\dagger - \hat{\mathbf{A}}_1^\dagger \hat{\mathbf{A}}_1 \hat{\rho} - \hat{\rho} \hat{\mathbf{A}}_1^\dagger \hat{\mathbf{A}}_1 \right) \\ & + \epsilon \left(2\hat{\mathbf{A}}_1^\dagger \hat{\rho} \hat{\mathbf{A}}_1 - \hat{\mathbf{A}}_1 \hat{\mathbf{A}}_1^\dagger \hat{\rho} - \hat{\rho} \hat{\mathbf{A}}_1 \hat{\mathbf{A}}_1^\dagger \right), \end{aligned} \quad (8)$$

where

$$\hat{\mathbf{H}}_0 = \hbar\omega \sum_{i=2}^N \hat{\mathbf{A}}_i^\dagger \hat{\mathbf{A}}_i + \hbar(\omega + \delta) \hat{\mathbf{A}}_1^\dagger \hat{\mathbf{A}}_1. \quad (9)$$

The real functions $\lambda, \delta, \epsilon$ are implicitly defined in terms of the auxiliary function $\eta(t)$

$$\eta(t) = \exp \left(- \int_0^t \lambda(t') dt' - i\omega t - i \int_0^t \delta(t') dt' \right), \quad (10)$$

which satisfies the integrodifferential equation

$$\dot{\eta} + i\omega\eta + \int_0^t d\tau \sum_k |c_k|^2 e^{i\omega_k(t-\tau)} \eta(\tau) = 0, \quad (11)$$

subject to the initial condition $\eta(0) = 1$. Moreover, considering the environment in thermal equilibrium, we have

$$\epsilon(t) = \frac{|\eta(t)|^2}{2} \frac{d}{dt} \left(\sum_k \frac{|c_k|^2 n_k(\beta)}{|\eta(t)|^2} \left| \int_0^t d\tau e^{-i\omega_k(t-\tau)} \eta(\tau) \right|^2 \right), \quad (12)$$

where $n_k(\beta)$ is the mean excitation number for the k^{th} mode of the environment at inverse temperature $\beta = 1/k_B T$. If the usual Born-Markov approximations hold, then $\delta(t) = 0$, $\lambda(t) = \sum_i k_i := k$, and $\epsilon = k\bar{n}$, where k_i characterize

the Markovian evolution when only the i^{th} original oscillator is coupled to the bath, and $\bar{n}$ is the environment mean number of thermal excitations at frequency ω . In this case the master equation simplifies to

$$\begin{aligned} \frac{d\hat{\rho}}{dt} = & -i\omega \sum_{i=1}^N \left[\hat{\mathbf{A}}_i^\dagger \hat{\mathbf{A}}_i, \hat{\rho} \right] + k(\bar{n} + 1) \left(2\hat{\mathbf{A}}_1 \hat{\rho} \hat{\mathbf{A}}_1^\dagger - \hat{\mathbf{A}}_1^\dagger \hat{\mathbf{A}}_1 \hat{\rho} - \hat{\rho} \hat{\mathbf{A}}_1^\dagger \hat{\mathbf{A}}_1 \right) \\ & + k\bar{n} \left(2\hat{\mathbf{A}}_1^\dagger \hat{\rho} \hat{\mathbf{A}}_1 - \hat{\mathbf{A}}_1 \hat{\mathbf{A}}_1^\dagger \hat{\rho} - \hat{\rho} \hat{\mathbf{A}}_1 \hat{\mathbf{A}}_1^\dagger \right). \end{aligned} \quad (13)$$

As one should expect by eq. (7), the master equation above describes the dissipative evolution of the collective mode $\hat{\mathbf{A}}_1$, and the independent unitary evolution of the remaining modes $\hat{\mathbf{A}}_i$, $i \geq 2$. It also should be noted that the damping constant k of mode $\hat{\mathbf{A}}_1$ is larger (in general, much larger, for large N) than the individual constants k_i of the modes $\hat{\mathbf{a}}_i$. This should be compared to the superradiance analogy discussed at section III.

V. TWO OSCILLATORS IN A DISSIPATIVE ENVIRONMENT

From now on, we specialize to the case of two harmonic oscillators ($N = 2$). The collective modes can thus be written as

$$\hat{\mathbf{A}}_1^\dagger = \cos(\theta)\hat{\mathbf{a}}_1^\dagger + \sin(\theta)\hat{\mathbf{a}}_2^\dagger, \quad \hat{\mathbf{A}}_2^\dagger = -\sin(\theta)\hat{\mathbf{a}}_1^\dagger + \cos(\theta)\hat{\mathbf{a}}_2^\dagger, \quad (14)$$

with $\tan \theta = G_2/G_1$ (this quotient can be taken as a positive number, if necessary, by redefining the mode $\hat{\mathbf{a}}_2$). If Markovian approximation is made, this relation takes the form $\tan \theta = \sqrt{k_2/k_1}$. The expression (14) can be considered as giving $\{\hat{\mathbf{A}}_i^\dagger\}$ by applying a rotation operator $\mathcal{R}(\theta)$ on the set $\{\hat{\mathbf{a}}_i^\dagger\}$. It is usual to call an operator which acts on operators a *superoperator*. Thus, $\mathcal{R}(\theta)$ is a *rotation superoperator*. It is convenient to represent superoperators using the algebraic relations among the operators in which they act on, and a useful notation is to introduce a dot ($\bullet$) in the position where the operator to be acted on must be placed. For example, $[\hat{\mathbf{A}}, \bullet] \hat{\mathbf{B}} = [\hat{\mathbf{A}}, \hat{\mathbf{B}}]$, and $[\hat{\mathbf{A}}, \bullet]^2 \hat{\mathbf{B}} = [\hat{\mathbf{A}}, [\hat{\mathbf{A}}, \hat{\mathbf{B}}]]$. With this convention in mind one can verify that

$$\mathcal{R}(\theta) = \exp \left\{ \theta [\hat{\mathbf{a}}_1 \hat{\mathbf{a}}_2^\dagger - \hat{\mathbf{a}}_2 \hat{\mathbf{a}}_1^\dagger, \bullet] \right\}. \quad (15)$$

Superoperators are very useful to study time evolution. As one can define an evolution operator $\hat{\mathbf{U}}(t)$ by $\hat{\mathbf{U}}(t)|\psi(0)\rangle = |\psi(t)\rangle$, the *evolution superoperator* is defined by $\mathcal{U}(t)\hat{\rho}(0) = \hat{\rho}(t)$. One completely solves the time evolution of a system by writing its evolution superoperator. The equation (8) can be solved by the superoperator

$$\mathcal{U}(t) = e^{-i\omega t [\hat{\mathbf{A}}_2^\dagger \hat{\mathbf{A}}_2, \bullet]} v e^{(1-v) \hat{\mathbf{A}}_1^\dagger \bullet \hat{\mathbf{A}}_1} e^{x \hat{\mathbf{A}}_1^\dagger \hat{\mathbf{A}}_1 \bullet} e^{x^* \bullet \hat{\mathbf{A}}_1^\dagger \hat{\mathbf{A}}_1} e^{z \hat{\mathbf{A}}_1 \bullet \hat{\mathbf{A}}_1^\dagger} \quad (16)$$

where the coefficients $v(t)$, $x(t)$ and $z(t)$ can be given in terms of the functions $\eta(t)$, eq. (10), and $\mathcal{N}(t)$,

$$\mathcal{N}(t) = \int_0^t d\tau \epsilon(\tau) \left| \frac{\eta(\tau)}{\eta(t)} \right|^2, \quad (17)$$

as follows

$$v(t) = \frac{1}{1 + \mathcal{N}(t)}, \quad x(t) = \ln \frac{\eta(t)}{\sqrt{1 + \mathcal{N}(t)}}, \quad z(t) = 1 - \frac{|\eta(t)|^{-2}}{1 + \mathcal{N}(t)}.$$

If the Markovian limit is applied, the preceding formulas reduce to

$$v = \frac{1}{1 + \bar{n}(1 - e^{-2(k_1+k_2)t})}, \quad x = \ln \frac{e^{(-i\omega - k_1 - k_2)t}}{\sqrt{1 + \bar{n}(1 - e^{-2(k_1+k_2)t})}}, \quad z = \frac{(\bar{n} + 1)(1 - e^{-2(k_1+k_2)t})}{1 + \bar{n}(1 - e^{-2(k_1+k_2)t})}. \quad (18)$$

The evolution superoperator $\mathcal{U}(t)$ can be expressed in terms of the original mode operators $\hat{\mathbf{a}}_i$ by using the rotation superoperator $\mathcal{R}(\theta)$ in the following way

$$\mathcal{U}(t) = e^{\theta [\hat{\mathbf{a}}_1 \hat{\mathbf{a}}_2^\dagger - \hat{\mathbf{a}}_2 \hat{\mathbf{a}}_1^\dagger, \bullet]} e^{-i\omega t [\hat{\mathbf{a}}_2^\dagger \hat{\mathbf{a}}_2, \bullet]} v e^{(1-v) \hat{\mathbf{A}}_1^\dagger \bullet \hat{\mathbf{A}}_1} e^{x \hat{\mathbf{A}}_1^\dagger \hat{\mathbf{A}}_1 \bullet} e^{x^* \bullet \hat{\mathbf{A}}_1^\dagger \hat{\mathbf{A}}_1} e^{z \hat{\mathbf{A}}_1 \bullet \hat{\mathbf{A}}_1^\dagger} e^{-\theta [\hat{\mathbf{a}}_1 \hat{\mathbf{a}}_2^\dagger - \hat{\mathbf{a}}_2 \hat{\mathbf{a}}_1^\dagger, \bullet]}. \quad (19)$$

Since the second collective mode is effectively decoupled from the environment, any density operator in the Hilbert space of this mode, times the asymptotic density operator of the coupled collective mode, provided it exists, will experience a unitary evolution. For simplicity we will further on restrict ourselves to the zero temperature case. Any density operator of the form (kets $|m, n\rangle$ refer to the original modes $\hat{\mathbf{a}}_i$):

$$\begin{aligned}\hat{\rho} &= \sum_{n,m} \frac{\rho_{n,m}}{n!m!} \left(\hat{\mathbf{A}}_2^\dagger\right)^n |0,0\rangle \langle 0,0| \left(\hat{\mathbf{A}}_2\right)^m \\ &= \sum_{n,m} \frac{\rho_{n,m}}{n!m!} \left(-\hat{\mathbf{a}}_1^\dagger \sin \theta + \hat{\mathbf{a}}_2^\dagger \cos \theta\right)^n |0,0\rangle \langle 0,0| \left(-\hat{\mathbf{a}}_1 \sin \theta + \hat{\mathbf{a}}_2 \cos \theta\right)^m \\ &= \sum_{n,m,n_1,m_1} \rho_{n,m} \frac{\sqrt{n!m!}(-\sin \theta)^{n+m-n_1-m_1}(\cos \theta)^{n_1+m_1}}{\sqrt{(n-n_1)!n_1!(m-m_1)!m_1!}} \\ &\quad \times |n_1, n-n_1\rangle \langle m_1, m-m_1|,\end{aligned}\tag{20}$$

will be protected against dissipation and decoherence. In fact, applying the evolution superoperator to an initial density matrix of this form, we obtain

$$\begin{aligned}\hat{\rho}(t) &= \sum_{n,m,n_1,m_1} e^{-i\omega t(n-m)} \rho_{n,m} \frac{\sqrt{n!m!}(-\sin \theta)^{n+m-n_1-m_1}(\cos \theta)^{n_1+m_1}}{\sqrt{(n-n_1)!n_1!(m-m_1)!m_1!}} \\ &\quad \times |n_1, n-n_1\rangle \langle m_1, m-m_1|,\end{aligned}\tag{21}$$

as one must expect.

Now, we use the evolution superoperator on the initial operator density

$$\hat{\rho}(0) = (\cos \alpha |1,0\rangle + e^{i\phi} \sin \alpha |0,1\rangle) (\cos \alpha \langle 1,0| + e^{-i\phi} \sin \alpha \langle 0,1|),\tag{22}$$

which can be viewed as a one photon Fock state of the mode given by the creation operator

$$\hat{\mathbf{A}}^\dagger(\alpha, \phi) = \cos \alpha \hat{\mathbf{a}}_1^\dagger + e^{i\phi} \sin \alpha \hat{\mathbf{a}}_2^\dagger,\tag{23}$$

where α and ϕ can be compared to Stokes parameters describing polarization. We will obtain how the dissipative properties of the mode (α, ϕ) depend on these parameters. As this is a natural way for experimentally test DFS[9], in the next section a more realistic situation is discussed. The above state asymptotically approaches the rank 2 density operator given by

$$\hat{\rho}_{t \rightarrow \infty} = P |\psi\rangle \langle \psi| + (1-P) |0,0\rangle \langle 0,0|\tag{24}$$

where the state $|\psi\rangle$ depends only on the individual decay rates k_i ,

$$|\psi\rangle = \frac{\sqrt{k_2} |1,0\rangle - \sqrt{k_1} |0,1\rangle}{\sqrt{k_1 + k_2}}\tag{25}$$

and the weight P of this state is given by

$$P = \langle \psi | \hat{\rho}(0) | \psi \rangle = \left| \frac{\sqrt{k_2} \cos \alpha - \sqrt{k_1} e^{i\phi} \sin \alpha}{\sqrt{k_1 + k_2}} \right|^2.\tag{26}$$

Observe that varying α and ϕ we can go from total preservation to total leakage. For example, if we set $\tan(\alpha) = \sqrt{k_1/k_2}$, and $\phi = 0$ then the full state will leak to the ground state $|0,0\rangle$, since in this case $\alpha = \theta$ and the initial photon was in the “superradiant” mode $\hat{\mathbf{A}}_1$. On the other hand, if we set $\tan(\alpha) = -\sqrt{k_2/k_1}$, and $\phi = 0$ then the initial state will be exactly equal to $|\psi\rangle$ (one photon in mode $\hat{\mathbf{A}}_2$), and will persist at all times with probability 1 (aside for an unimportant global phase). All other combinations will go to the density operator (24), which can be considered as an ensemble of pure state $|\psi\rangle$ with probability P and the ground state $|0,0\rangle$ with probability $1-P$. One can define the asymptotic fidelity, $F_\infty(\alpha, \phi)$, which is the overlap between the initial and asymptotic density matrices. In the above example it is given by

$$F_\infty(\alpha, \phi) = \left| \frac{(\sqrt{k_2} \cos(\alpha) - \sqrt{k_1} e^{i\phi} \sin(\alpha))(\sqrt{k_2} \cos(\alpha) - \sqrt{k_1} e^{-i\phi} \sin(\alpha))}{k_1 + k_2} \right|^2.\tag{27}$$

VI. EFFECTS OF MORE REALISTIC MODELING

We remark that the results above were obtained under a number of assumptions, which will be relaxed below. Notice that the use of the rotating wave approximation (RWA) is not essential in obtaining the decoupled mode: any interaction linear in the field operators would be as good (provided the other assumptions hold). Had we chosen an interaction linear in the identical oscillators but nonlinear on the environmental operators we would have obtained also a decoupled collective mode. In these cases, however, the complication would be only of technical nature leading to (much) more complex dynamics.

Another important hypothesis to obtain DFS is that of identical frequencies of the original main oscillators. Of course, any interaction between them would destroy the symmetry upon which the existence of DFS rests. On the other hand, we have assumed that the oscillator-environment coupling satisfies $g_{ik} = G_i D_k$, which amounts to a separable coupling. It is not an easy task to find realizations of such interactions in nature given its nonlocal character. However, it might be a good approximation in special circumstances, as *e.g.* optical cavities. A particular consequence of the separability hypothesis can be seen writing the master equation, in the zero temperature limit, in terms of the original oscillators (with different frequencies for generality)

$$\begin{aligned} \mathcal{L}_0 = & (-i\omega_1 - k_1) \hat{\mathbf{a}}_1^\dagger \hat{\mathbf{a}}_1 \bullet + (i\omega_1 - k_1) \bullet \hat{\mathbf{a}}_1^\dagger \hat{\mathbf{a}}_1 + 2k_1 \hat{\mathbf{a}}_1 \bullet \hat{\mathbf{a}}_1^\dagger + \\ & (-i\omega_2 - k_2) \hat{\mathbf{a}}_2^\dagger \hat{\mathbf{a}}_2 \bullet + (i\omega_2 - k_2) \bullet \hat{\mathbf{a}}_2^\dagger \hat{\mathbf{a}}_2 + 2k_2 \hat{\mathbf{a}}_2 \bullet \hat{\mathbf{a}}_2^\dagger + \\ & k_3 \left(2\hat{\mathbf{a}}_1 \bullet \hat{\mathbf{a}}_2^\dagger - \hat{\mathbf{a}}_2^\dagger \hat{\mathbf{a}}_1 \bullet - \bullet \hat{\mathbf{a}}_2^\dagger \hat{\mathbf{a}}_1 \right) + k_3^* \left(2\hat{\mathbf{a}}_2 \bullet \hat{\mathbf{a}}_1^\dagger - \hat{\mathbf{a}}_1^\dagger \hat{\mathbf{a}}_2 \bullet - \bullet \hat{\mathbf{a}}_1^\dagger \hat{\mathbf{a}}_2 \right). \end{aligned} \quad (28)$$

The new quantity k_3 appears since we consider the same environment interacting with both oscillators. These terms can be considered as an interaction between the oscillators mediated by the environment. If the separability condition is fulfilled, $|k_3|^2 = k_1 k_2$, and if the oscillators are identical, eq. (28) is the same as eq. (13). The other limit case is to consider both oscillators interacting independently with the environment. In this situation, the independence of the phases of the interaction coefficients g_{ik} will make their net effect on k_3 null, and the eq. (28) will just describe two independent damped harmonic oscillators. Our interest is to study the above equation when the conditions for the existence of DFS are almost satisfied, *i.e.* $|k_3|^2 \approx k_1 k_2$ and $\omega_1 \approx \omega_2$. It must be noted that, by Cauchy-Schwarz inequality, $|k_3|^2 \leq k_1 k_2$.

We should point out that we are not using the usual approach of perturbation theory, of adding a small perturbation $\varepsilon \hat{\mathbf{H}}'$. In Ref. [15], the authors have shown that DFS are robust up to order ε in the perturbation, and all orders in time, so our approach must be connected to second order (in ε) perturbation.

The explicit solution to this problem is given in the appendix. As is expected, there is no DFS without the separability and degeneracy assumptions, but if we are close to this conditions, we can obtain states much more robust to decoherence and dissipation than others. As in the previous section, consider the one photon states of eq. (22). Then we can define a *weak decoherence mode* (WD), which tends to the DFS when degeneracy and separability are approximated, and a *strong decoherence mode* (SD) which is analogous to the superradiant mode $\hat{\mathbf{A}}_1$. We want to explore the slight deviations from separability and degeneracy, so we define δk and $\delta\omega$ by

$$\delta k = \sqrt{k_1 k_2} - |k_3|, \quad 2\delta\omega = \omega_2 - \omega_1, \quad (29)$$

and consider $\delta\omega \ll \omega_i$ and $\delta k \ll k_i$.

As in the previous section, varying the parameters of the initial state (22) can be interpreted as varying the mode of the initial photon. In the regime discussed above we obtain:

$$\hat{\mathbf{A}}_{\text{SD}}^\dagger = \frac{1}{\sqrt{k_1 + k_2}} \left(\sqrt{k_1} \hat{\mathbf{a}}_1^\dagger \pm \sqrt{k_2} e^{\pm i \frac{\delta\omega}{k}} \hat{\mathbf{a}}_2^\dagger \right), \quad (30)$$

which must be compared to eq. (23). The new parameter k is a kind of effective mean damping, and in the regime here discussed can be considered as $k \approx (k_1 + k_2)/2$. Each mode has its own damping constant, and this two are the extrema. Explicitly we have

$$k_{\text{WD}} = \frac{2\delta k \sqrt{k_1 k_2}}{k_1 + k_2} \approx \delta k, \quad (31)$$

for the weak decoherence mode, and

$$k_{\text{SD}} = k_1 + k_2 \approx 2k \quad (32)$$

for the strong decoherence mode. One must note that while k_{SD} is of the same order as the individual damping constants k_i , the value of k_{WD} can be much lower. In the experiment with ions[9] it was exactly this lowering of the damping constant that was exhibited as an evidence of decoherence “free” subspaces. In the same experiment, one can see that the difference in damping constants is much larger in the situation with an engineered noise applied, since in this case the interaction with the environment is much closer to the separability condition.

VII. CONCLUDING REMARKS

We have studied decoherence and dissipation free modes for systems of harmonic oscillators. We discussed sufficient conditions for their existence. Although theoretically simple, these conditions are very difficult to be implemented in practical experiments. So we studied the slight deviations of this conditions, and instead of decoherence free subspaces, we obtained *weak decoherence modes*. This suggests that weak decoherence subspaces can be used to store quantum information for times much larger than individual carriers would be able to, even without being rigorous DFS.

We compare DFS to the so called super and subradiance effects of a maser. In fact, when in the last section we compare the damping constants for weak and strong decoherence (eqs. (31) and (32)), this effect mimimimics interference problem, where we are comparing the maximum and the minimum of a certain quantity in which interference effects are recorded (in this case, the damping constant).

It is important to stress that larger deviations of the rigorous conditions for DFS preclude the existence of even weakly decoherence subspaces, by making the decoherence time scales for such states smaller. However, one can conjecture that this kind of mechanism is so general that whenever an experiment obtain quantum mechanical results, it is testing some kind of DFS (*e.g.* the fullerenes experiment[22]).

Acknowledgments

We gratefully acknowledge comments from A. N. Salgueiro. This work was partly funded by FAPESP, CNPq and PRONEX (Brazil), and Colciencias, DINAIN (Colombia). K.M.F.R. gratefully acknowledges the Instituto de Física, Universidade de São Paulo, for their hospitality and PRONEX for partial support.

Appendix: Realistic model of two oscillators in details

The evolution superoperator for eq.(28) can be expressed as[21]

$$\begin{aligned} \mathcal{U}(t) = & e^{j_1(t)\hat{\mathbf{a}}_1\cdot\hat{\mathbf{a}}_1^\dagger} e^{j_2(t)\hat{\mathbf{a}}_2\cdot\hat{\mathbf{a}}_2^\dagger} e^{z(t)\hat{\mathbf{a}}_2\cdot\hat{\mathbf{a}}_1^\dagger} e^{z^*(t)\hat{\mathbf{a}}_1\cdot\hat{\mathbf{a}}_2^\dagger} e^{q(t)\hat{\mathbf{a}}_1\hat{\mathbf{a}}_2^\dagger} e^{q^*(t)\cdot\hat{\mathbf{a}}_1^\dagger\hat{\mathbf{a}}_2} e^{m_2(t)\hat{\mathbf{a}}_2^\dagger\hat{\mathbf{a}}_2} e^{m_2^*(t)\cdot\hat{\mathbf{a}}_2^\dagger\hat{\mathbf{a}}_2} \odot \\ & \odot e^{m_1(t)\hat{\mathbf{a}}_1^\dagger\hat{\mathbf{a}}_1} e^{m_1^*(t)\cdot\hat{\mathbf{a}}_1^\dagger\hat{\mathbf{a}}_1} e^{q(t)\hat{\mathbf{a}}_1^\dagger\hat{\mathbf{a}}_2} e^{q^*(t)\cdot\hat{\mathbf{a}}_1^\dagger\hat{\mathbf{a}}_2} \end{aligned} \quad (33)$$

where

$$R = \frac{k_2 + k_1}{2} + \frac{i(\omega_2 + \omega_1)}{2}, \quad c = k_2 - k_1 + i(\omega_2 - \omega_1), \quad r = \sqrt{c^2 + 4k_3^2}, \quad \Delta_{\pm} = c \pm r \quad (34)$$

$$q(t) = 2k_3 (1 - e^{r t}) (\Delta_+ e^{r t} - \Delta_-)^{-1} \quad \text{for } r \neq 0 \quad (35)$$

$$e^{m_1(t)} = \frac{e^{-R t}}{2r} e^{-\frac{r t}{2}} (\Delta_+ e^{r t} - \Delta_-), \quad e^{m_2(t)} = e^{-2R t} e^{-m_1(t)} \quad (36)$$

$$j_2(t) = (1 + |q(t)|^2) \left(\left| e^{m_2(t)} \right|^{-2} \right) - 1 \quad (37)$$

$$j_1(t) = \left| e^{-m_1(t)} + q(t)^2 e^{-m_2(t)} \right|^2 + (|q(t)|^2) \left(\left| e^{m_2(t)} \right|^{-2} \right) - 1 \quad (38)$$

$$z(t) = -q(t) e^{-(m_1^*(t) + m_2(t))} - q^*(t) (1 + |q(t)|^2) \left| e^{m_2(t)} \right|^{-2}. \quad (39)$$

In this calculation neither the separability nor the degeneracy (even approximated) conditions have been used so far. For the sake of comparison we use the same initial condition of section V (eq. (22)). In the general case its time evolution is also given by

$$\hat{\rho}(t) = P(t) |\psi(t)\rangle \langle \psi(t)| + (1 - P(t)) |0, 0\rangle \langle 0, 0| \quad (40)$$

but now the state $|\psi(t)\rangle$ is,

$$|\psi(t)\rangle = \frac{(\cos(\theta)M_-(t) + \sin(\theta)e^{i\phi}Q(t))|1,0\rangle + (\sin(\theta)e^{i\phi}M_+(t) + \cos(\theta)Q(t))|0,1\rangle}{\sqrt{P(t)}}, \quad (41)$$

and its coefficient is given by

$$P(t) = |\cos(\theta)M_-(t) + \sin(\theta)e^{i\phi}Q(t)|^2 + |\sin(\theta)e^{i\phi}M_+(t) + \cos(\theta)Q(t)|^2, \quad (42)$$

where the functions $M_{\pm}(t)$ and $Q(t)$ are given by

$$M_{\pm}(t) = \frac{e^{-Rt}}{2} \left(e^{-rt/2} \left(1 \mp \frac{c}{r} \right) + e^{rt/2} \left(1 \pm \frac{c}{r} \right) \right), \quad Q(t) = \frac{k_3}{r} e^{-Rt} \left(e^{-rt/2} - e^{rt/2} \right). \quad (43)$$

Now, we assume slight deviations from degeneracy and separability, that is, $\omega_1 = \omega - \delta\omega$, $\omega_2 = \omega + \delta\omega$, $k_3 = \sqrt{k_1 k_2} - \delta k$, with $\delta\omega \ll \omega$, $\delta k \ll \sqrt{k_1 k_2}$. Then, the state $|\psi(t)\rangle$ can be approximated as

$$|\psi(t)\rangle = e^{-i\omega t} \left(\frac{\zeta_1 |1,0\rangle + \xi_1 |0,1\rangle}{\sqrt{P(t)}} e^{-(k_1+k_2)t} + \frac{\zeta_2 |1,0\rangle + \xi_2 |0,1\rangle}{\sqrt{P(t)}} e^{-\frac{2\delta k \sqrt{k_1 k_2}}{k_1+k_2} t} \right), \quad (44)$$

where ζ_i, ξ_i do not have any temporal dependence and are given by

$$\zeta_1 = \frac{(k_1 \mp i\delta\omega) \cos(\alpha) \pm \sqrt{k_1 k_2} \sin(\alpha) e^{i\phi}}{k_1 + k_2}, \quad (45)$$

$$\xi_1 = \frac{(k_2 \pm i\delta\omega) e^{i\phi} \sin(\alpha) \pm \sqrt{k_1 k_2} \cos(\alpha)}{k_1 + k_2}. \quad (46)$$

In the general case it is not possible to find initial conditions which are completely decoherence free. Nor it is possible to find two orthogonal subspaces with very different characters in what decoherence is concerned. However, we can choose the initial condition as to have a minimal component either in a strong decoherence (SD) or in a weak decoherence (WD) subspaces, by choosing, e.g.

$$\tan(\alpha)_{\text{WD}}^{\text{SD}} = \pm \sqrt{\frac{k_2}{k_1}}, \quad \phi_{\text{WD}}^{\text{SD}} = \pm \frac{\delta\omega}{k}, \quad (47)$$

where k is some average dissipation constant. The corresponding states, apart from a phase, can be written as

$$|\psi_{\text{WD}}^{\text{SD}}\rangle = \frac{1}{\sqrt{k_1 + k_2}} \left(\sqrt{k_1} |1,0\rangle \pm \sqrt{k_2} e^{\pm i\delta\omega/k} |0,1\rangle \right). \quad (48)$$

If $\delta\omega \ll k_1, k_2$ the phase can be ignored. Moreover, if $k_1 = k_2$ then we have $k = k_1 = k_2$ and the phase can be unambiguously determined. The weak decoherence wavefunction defines a mode which is robust against decoherence. The damping constant of this mode can be read from eq. (44) as the value given in eq. (31). Analogously for the strong decoherence mode, with damping constant given by eq. (32).

-
- [1] M. A. Nielsen and I. L. Chuang, *Quantum computation and quantum information* (Cambridge University Press, Cambridge, 2000).
 - [2] W. H. Zurek, *Physics Today* **44** (Oct.), 36 (1991). D. Giulini *et al.*, *Decoherence and the appearance of a Classical World in Quantum Theory*, (Springer, 1996).
 - [3] P. W. Shor, Phys. Rev. A **52**, 2493 (1995); A. Ekert and C. Macchiavello, Phys. Rev. Lett. **77**, 2585 (1996); D. Gottesman, Phys. Rev. A **54**, 1862 (1996); A. R. Calderbank *et al.*, Phys. Rev. Lett. **78**, 405 (1997).
 - [4] L. Viola, E. Knill, and S. Lloyd. Phys. Rev. Lett. **82**, 2417 (1999). D. Vitali, and P. Tombesi. Phys. Rev. A **65**, 012305 (2001).
 - [5] G. M. Palma, K.-A. Suominen, A. K. Ekert. *Proc. R. Soc. London Ser. A*, **452**, 567 (1996); L.-M. Duan, G.-C. Guo, Phys. Rev. Lett. **79**, 1953 (1997); A. Beige, D. Braun, B. Tregenna, P. L. Knight. Phys. Rev. Lett. **85**, 1762 (2000).

- [6] D. A. Lidar, I. L. Chuang, K. B. Whaley, Phys. Rev. Lett. **81**, 2594 (1998).
- [7] P. Zanardi, M. Rasetti. Phys. Rev. Lett. **79**, 3306 (1997); P. Zanardi, Phys. Rev. A **60**, R729 (1999).
- [8] P. G. Kwiat *et al.*, *Science*, **290**, 498 (2000).
- [9] D. Kielpinski *et al.*, *Science*, **291**, 1013 (2001).
- [10] Q. A. Turchette *et al.*, Phys. Rev. Lett. **75**, 4710 (1995); A. Imamoglu, *et al.*, *ibid.*, **83**, 4204 (1999); A. Rauschenbeutel *et al.*, *ibid.*, **83**, 5166 (1999).
- [11] J. I. Cirac, and P. Zoller. *Nature*, **404**, 579, (2000), Phys. Rev. Lett. **74**, 4091 (1995); C. Monroe *et al.*, *ibid.*, **75**, 4714 (1995); K. Molmer, and A. Sorensen, *ibid.*, **82**, 1835 (1999); C. A. Sackett *et al.*, *Nature*, **393**, 133, (2000).
- [12] N. A. Garshenfeld, and I. L. Chuang, *Science*, **275**, 350 (1997); E. Knill *et al.*, Phys. Rev. A **57**, 3348 (1998); J. A. Jones *et al.*, *Nature*, **393**, 344, (1998); B. E. Kane, *Nature*, **393**, 133, (1998).
- [13] A. Barenco *et al.*, Phys. Rev. Lett. **74**, 4083 (1995); D. Loss, and D.P. DiVincenzo, Phys. Rev. A **57** 120 (1998); G. Burkard *et al.*, Phys. Rev. B **59**, 2070 (1999); L. Quiroga and N.F. Johnson, Phys. Rev. Lett. **83**, 2270 (1999); J.H. Reina *et al.*, Phys. Rev. A **62**, 12305 (2000); J.H. Reina *et al.*, Phys. Rev. B **62**, R2267 (2000), F. Troiani *et al.*, Phys. Rev. B **62**, R2263 (2000); E. Biolatti *et al.*, Phys. Rev. Lett. **85**, 5647 (2000).
- [14] L.-A. Wu, D. A. Lidar. Phys. Rev. Lett. **88**, 207902 (2002).
- [15] D. Bacon, D. A. Lidar, K. B. Whaley, Phys. Rev. A **60**, 1944 (1999).
- [16] As must be expected, the experimental verification could not find a completely isolated subspace as the simple theory foresee, but just a sensible reduction on decoherence rate as compared to other subspaces. This can also be viewed as a motivation to our work: to study the effects of small deviations on the ideal situation.
- [17] L. Davidovich *et al.*, Phys. Rev. Lett. **71**, 2360 (1993).
- [18] R. H. Dicke, *Phys. Rev.* **93**, 99 (1954).
- [19] K. M. Fonseca-Romero and M. C. Nemes, Report No. quant-ph/0201107.
- [20] R. P. Feynman, R. B. Leighton, and M. Sands, *The Feynman Lectures on Physics* vol 3, Addison-Wesley (1963).
- [21] S. G. Mokarzel, *Ph.D. Thesis*, State University of São Paulo (2000).
- [22] M. Arndt *et al.*, *Nature* **401**, 680 (1999).